

Multibracket simple Lie algebras *

J.A. de Azcárraga[†] and J.C. Pérez Bueno[‡]

Department of Applied Mathematics and Theoretical Physics, Silver St.,
Cambridge CB3 9EW, UK

Abstract

We introduce higher-order (or multibracket) simple Lie algebras that generalize the ordinary Lie algebras. Their ‘structure constants’ are given by Lie algebra cohomology cocycles which, by virtue of being such, satisfy a suitable generalization of the Jacobi identity. Finally, we introduce a nilpotent, complete BRST operator associated with the l multibracket algebras which are based on a given simple Lie algebra of rank l .

Given $[X, Y] := XY - YX$, the standard Jacobi identity (JI) $[[X, Y], Z] + [[Y, Z], X] + [[Z, X], Y] = 0$ is automatically satisfied if the product is associative. For a Lie algebra \mathcal{G} , $[X_i, X_j] = C_{ij}^k X_k$, the JI may be written in terms of C_{ij}^k as

$$\frac{1}{2} \epsilon_{i_1 i_2 i_3}^{j_1 j_2 j_3} C_{j_1 j_2}^\rho C_{\rho j_3}^\sigma = 0 \quad . \quad (1)$$

Let \mathcal{G} be simple and (for simplicity) compact. Then, the Killing metric k , with coordinates $k_{ij} = k(X_i, X_j)$, is non-degenerate and, after suitable normalization, can be brought to the form $k_{ij} = \delta_{ij}$. Moreover, k is an invariant polynomial, *i.e.*

$$k([Y, X], Z) + k(X, [Y, Z]) = 0 \quad . \quad (2)$$

We also know that k defines the second order Casimir invariant. Using this symmetric polynomial we may always construct a non-trivial three-cocycle

$$\omega_{i_1 i_2 i_3} := k([X_{i_1}, X_{i_2}], X_{i_3}) = C_{i_1 i_2}^\rho k_{\rho i_3} \quad (3)$$

which is indeed skew-symmetric as consequence of (1) or (2).

In fact, it is known since the classical work of Cartan, Pontrjagin, Hopf and others that, from a topological point of view, the group manifolds of all simple compact groups are essentially equivalent to (have the [real] homology of) products of odd

*Talk at XXI Int. Coll. on Group Theor. Methods in Phys. (Goslar, July 1996), to appear in the Proceedings.

[†]St. John’s College Overseas Visiting Scholar.

[‡]On sabbatical (J.A.) leave and on leave of absence (J.C.P.B.) from Departamento de Física Teórica and IFIC (Centro Mixto Univ. de Valencia-CSIC), E-46100 Burjassot (Valencia), Spain.

spheres, that S^3 is always present in these products and that the simple Lie algebra cocycles are, via the ‘localization’ process, in one-to-one correspondence with bi-invariant de Rham cocycles on the associated compact group manifolds G . This is due to the intimate relation between the order of the $l(=\text{rank } \mathcal{G})$ primitive symmetric polynomials which can be defined on a simple Lie algebra, their l associated generalized Casimir-Racah invariants [1] and the topology of the associated simple groups. Such a relation was a key fact in the eighties for the understanding of non-abelian anomalies in gauge theories [2].

The simplest (of order 3) higher-order invariant polynomial $d_{ijk} = d(X_i, X_j, X_k)$ appears for $su(3)$ (and only for A_l -type algebras, $l \geq 2$); it is given by the symmetric trace of three $su(3)$ generators. From d_{ijk} we may construct

$$\omega_{i_1 i_2 i_3 i_4 i_5} := \epsilon_{i_2 i_3 i_4}^{j_2 j_3 j_4} d([X_{i_1}, X_{j_2}], [X_{j_3}, X_{j_4}], X_{i_5}) = \epsilon_{i_2 i_3 i_4}^{j_2 j_3 j_4} C_{i_1 j_2}^\rho C_{j_3 j_4}^\sigma d_{\rho \sigma i_5} \quad (4)$$

(cf. (3)), and it can be checked that (4) defines a fifth-order invariant form (the proof will be given in the general case). The existence of this five-form ω shows us that $su(3)$ is, from a topological point of view, equivalent to $S^3 \times S^5$. If we calculate in $su(3)$ the ‘four-bracket’ we find that

$$[X_{j_1}, X_{j_2}, X_{j_3}, X_{j_4}] = \sum_{s \in S_4} \pi(s) X_{s(j_1)} X_{s(j_2)} X_{s(j_3)} X_{s(j_4)} = \omega_{j_1 j_2 j_3 j_4}^\sigma X_\sigma \quad , \quad (5)$$

where the generators X_i may be taken proportional to the Gell-Mann matrices, $X_i = \frac{\lambda_i}{2}$, and $\pi(s)$ is the parity sign of the permutation s . Thus, $\omega_{j_1 j_2 j_3 j_4}^\sigma$ is related to the four-bracket and a five-cocycle (five-form) in the same way as $C_{j_1 j_2}^\sigma$ is associated with the standard Lie bracket and a three-cocycle (three-form).

We may ask ourselves whether this construction could be extended to all the higher-order polynomials to define from them higher-order simple Lie algebras satisfying an appropriate generalization of the JI. The affirmative answer is given in [3]; we outline below the main steps that led to it. It is interesting to note that this construction may be used to produce examples of a generalization [4] of the Poisson structure different from that underlying Nambu mechanics [5].

a) Invariant polynomials on the Lie algebra \mathcal{G}

Let T_i be the elements of a representation of \mathcal{G} . Then the symmetric trace $k_{i_1 \dots i_m} \equiv \text{sTr}(T_{i_1} \dots T_{i_m})$ (we shall only consider here sTr although not all invariant polynomials are of this form [1]; see [3]) verifies the invariance condition

$$\sum_{s=1}^m C_{\nu i_s}^\rho k_{i_1 \dots i_{s-1} \rho i_{s+1} \dots i_m} = 0 \quad . \quad (6)$$

Proof: By definition of k , the *l.h.s.* of (6) (cf. (2)) is

$$\text{sTr} \left(\sum_{s=1}^m T_{i_1} \dots T_{i_{s-1}} [T_\nu, T_{i_s}] T_{i_{s+1}} \dots T_{i_m} \right) = \text{sTr} (T_\nu T_{i_1} \dots T_{i_m} - T_{i_1} \dots T_{i_m} T_\nu) = 0 \quad , \quad (7)$$

q.e.d. The above symmetric polynomial is associated to an invariant symmetric tensor field on the group G associated with \mathcal{G} , $k(g) = k_{i_1 \dots i_m} \omega^{i_1}(g) \otimes \dots \otimes \omega^{i_m}(g)$, where the $\omega^i(g)$ are left invariant one-forms on G . Since the Lie derivative of ω^k is given by $L_{X_i} \omega^k = -C_{ij}^k \omega^j$ for a LI vector field X_i on G , the invariance condition is the statement

$$(L_{X_\nu} k)(X_{i_1}, \dots, X_{i_m}) = - \sum_{s=1}^m k(X_{i_1}, \dots, [X_\nu, X_{i_s}], \dots, X_{i_m}) = 0 \quad (8)$$

c.f. (2). On forms, the invariance condition (8) may be written as

$$\epsilon_{i_1 \dots i_q}^{j_1 \dots j_q} C_{\nu j_1}^\rho \omega_{\rho j_2 \dots j_q} = 0 \quad . \quad (9)$$

b) Invariant forms on the Lie group G

Let $k_{i_1 \dots i_m}$ be an invariant symmetric polynomial on \mathcal{G} and let us define

$$\tilde{\omega}_{\rho j_2 \dots j_{2m-2} \sigma} := k_{i_1 \dots i_{m-1} \sigma} C_{\rho j_2}^{i_1} \dots C_{j_{2m-3} j_{2m-2}}^{i_{m-1}} \quad . \quad (10)$$

Then the odd order $(2m-1)$ -tensor

$$\omega_{\rho l_2 \dots l_{2m-2} \sigma} := \epsilon_{l_2 \dots l_{2m-2}}^{j_2 \dots j_{2m-2}} \tilde{\omega}_{\rho j_2 \dots j_{2m-2} \sigma} \quad (11)$$

is a fully skew-symmetric tensor. We refer to Lemma 8.1 in [4] for the proof.

Moreover, ω is an invariant form because for $q = 2m-1$ the *l.h.s.* of (9) is

$$\begin{aligned} \epsilon_{i_1 \dots i_{2m-1}}^{j_1 \dots j_{2m-1}} C_{\nu j_1}^\rho \omega_{j_2 \dots j_{2m-1} \rho} &= \epsilon_{i_1 \dots i_{2m-1}}^{j_1 \dots j_{2m-1}} C_{\nu j_1}^\rho \epsilon_{j_3 \dots j_{2m-1}}^{l_3 \dots l_{2m-1}} \tilde{\omega}_{j_2 l_3 \dots l_{2m-1} \rho} \\ &= (2m-3)! \epsilon_{i_1 \dots i_{2m-1}}^{j_1 \dots j_{2m-1}} k_{l_1 \dots l_m} C_{\nu j_1}^{l_1} \dots C_{j_{2m-2} j_{2m-1}}^{l_m} \\ &= (2m-3)! \epsilon_{i_1 \dots i_{2m-1}}^{j_1 \dots j_{2m-1}} \left[\sum_{s=2}^m k_{\nu l_2 \dots l_{s-1} \rho l_{s+1} \dots l_m} C_{j_1 l_s}^\rho \right] C_{j_2 j_3}^{l_2} \dots C_{j_{2m-2} j_{2m-1}}^{l_m} = 0 \quad . \end{aligned} \quad (12)$$

This result follows recalling

$$\epsilon_{i_1 \dots i_p i_{p+1} \dots i_n}^{j_1 \dots j_p j_{p+1} \dots j_n} \epsilon_{j_{p+1} \dots j_n}^{l_{p+1} \dots l_n} = (n-p)! \epsilon_{i_1 \dots i_p i_{p+1} \dots i_n}^{j_1 \dots j_p l_{p+1} \dots l_n} \quad (13)$$

in the second equality, using the invariance of k [eq. (6)] in the third one and the JI in the last equality for each of the $(m-1)$ terms in the bracket.

This may be seen without using coordinates; indeed (10) is expressed as

$$\tilde{\omega}(X_\rho, X_{j_2}, \dots, X_{j_{2m-2}}, X_\sigma) := k([X_\rho, X_{j_2}], \dots, [X_{j_{2m-3}}, X_{j_{2m-2}}], X_\sigma) \quad , \quad (14)$$

and the $(2m-1)$ -form ω is obtained antisymmetrizing (14) as in (11) (cf. (4)). Hence

$$(L_{X_\nu} \tilde{\omega})(X_{i_1}, \dots, X_{i_{2m-1}}) = - \sum_{p=1}^{2m-1} \tilde{\omega}(X_{i_1}, \dots, [X_\nu, X_{i_p}], \dots, X_{i_{2m-1}})$$

$$\begin{aligned}
&= - \sum_{s=1}^{m-1} k([X_{i_1}, X_{i_2}], \dots, [[X_\nu, X_{i_{2s-1}}], X_{i_{2s}}] + [X_{i_{2s-1}}, [X_\nu, X_{i_{2s}}]], \dots, \\
&\quad [X_{i_{2m-3}}, X_{i_{2m-2}}], X_{i_{2m-1}}) - k([X_{i_1}, X_{i_2}], \dots, [X_{i_{2m-3}}, X_{i_{2m-2}}], [X_\nu, X_{i_{2m-1}}]) \\
&= - \sum_{s=1}^{m-1} k([X_{i_1}, X_{i_2}], \dots, [X_\nu, [X_{i_{2s-1}}, X_{i_{2s}}]], \dots, [X_{i_{2m-3}}, X_{i_{2m-2}}], X_{i_{2m-1}}) \\
&\quad - k([X_{i_1}, X_{i_2}], \dots, [X_{i_{2m-3}}, X_{i_{2m-2}}], [X_\nu, X_{i_{2m-1}}]) \\
&= (L_{X_\nu} k)([X_{i_1}, X_{i_2}], \dots, [X_{i_{2m-3}}, X_{i_{2m-2}}], X_{i_{2m-1}}) = 0 \quad ; \tag{15}
\end{aligned}$$

where the JI has been used in the third equality and (8) in the last, *q.e.d.*

c) The generalized Jacobi condition

Now we are ready to check that the tensor ω introduced above verifies a generalized Jacobi condition that extends eq. (1) to multibracket algebras.

Theorem Let \mathcal{G} be a simple compact algebra, and let ω be the non-trivial Lie algebra $(2p+1)$ -cocycle obtained from the associated p invariant symmetric tensor on \mathcal{G} . Then ω verifies the *generalized Jacobi condition* (GJC)

$$\epsilon_{i_1 \dots i_{4p-1}}^{j_1 \dots j_{4p-1}} \omega_{\sigma j_1 \dots j_{2p-1}} \cdot^\rho \omega_{\rho j_{2p} \dots j_{4p-1}} = 0 \quad . \tag{16}$$

Proof: Using (11), (10) and (13), the *l.h.s.* of (16) is equal to

$$\begin{aligned}
&- (2p-3)! \epsilon_{i_1 \dots i_{4p-1}}^{j_1 \dots j_{4p-1}} k_{l_1 \dots l_p \sigma} C_{\rho j_1}^{l_1} \dots C_{j_{2p-2} j_{2p-1}}^{l_p} \omega_{j_{2p} \dots j_{4p-1}}^\rho \\
&= - (2p-3)! \epsilon_{i_1 \dots i_{4p-1}}^{j_1 \dots j_{4p-1}} k_{\dots l_p \sigma}^{l_1} C_{j_2 j_3}^{l_2} \dots C_{j_{2p-2} j_{2p-1}}^{l_p} C_{l_1 j_1}^\rho \omega_{\rho j_{2p} \dots j_{4p-1}} = 0 \quad , \tag{17}
\end{aligned}$$

where the invariance of ω (eq. (9)) has been used in the last equality, *q.e.d.*

d) Multibrackets and higher-order simple Lie algebras

Eq. (16) now allows us to define higher-order simple Lie algebras based on \mathcal{G} using [3] the Lie algebra cocycles ω on \mathcal{G} as generalized structure constants:

$$[X_{i_1}, \dots, X_{i_{2m-2}}] = \omega_{i_1 \dots i_{2m-2}}^\sigma X_\sigma \quad . \tag{18}$$

The GJC (16) satisfied by the cocycles is necessary since for *even* n -brackets of associative operators one has the generalized Jacobi identity

$$\frac{1}{(n-1)!n!} \sum_{\sigma \in S_{2n-1}} (-)^{\pi(\sigma)} [[X_{\sigma(1)}, \dots, X_{\sigma(n)}], X_{\sigma(n+1)}, \dots, X_{\sigma(2n-1)}] = 0 \quad . \tag{19}$$

This establishes the link between the \mathcal{G} -based *even* multibracket algebras and the *odd* Lie algebra cohomology cocycles on \mathcal{G} (note that for n odd the *l.h.s* is proportional to the odd $(2n-1)$ -multibracket $[X_1, \dots, X_{2n-1}]$ [3]).

Finally we comment that just in the same way that we can introduce for a Lie algebra a BRST nilpotent operator by

$$s = -\frac{1}{2} c^i c^j C_{ij}^k \frac{\partial}{\partial c^k} \equiv s_2 \quad , \quad s^2 = 0 \quad , \tag{20}$$

with $c^i c^j = -c^j c^i$, the set of invariant forms ω associated with a simple \mathcal{G} allows us to *complete* this operator in the form

$$s = -\frac{1}{2}c^{j_1}c^{j_2}\omega_{j_1j_2}{}^\sigma \frac{\partial}{\partial c^\sigma} - \dots - \frac{1}{(2m_i - 2)!}c^{j_1} \dots c^{j_{2m_i-2}}\omega_{j_1 \dots j_{2m_i-2}}{}^\sigma \frac{\partial}{\partial c^\sigma} - \dots$$

$$-\frac{1}{(2m_l - 2)!}c^{j_1} \dots c^{j_{2m_l-2}}\omega_{j_1 \dots j_{2m_l-2}}{}^\sigma \frac{\partial}{\partial c^\sigma} \equiv s_2 + \dots + s_{2m_i-2} + \dots + s_{2m_l-2}. \quad (21)$$

This new nilpotent operator s is the *complete BRST operator* [3] associated with \mathcal{G} .

For the relation of these constructions with the strongly homotopy algebras [6], possible extensions and connections with physics we refer to [3] and references therein.

Acknowledgements

This research has been partially supported by the CICYT and DGICYT, Spain (AEN 96-1669, PR 95-439). Both authors wish to thank the kind hospitality extended to them at DAMTP. Finally, the support of St. John's College (J.A.) and an FPI grant from the Spanish M.E.C. and the CSIC (J.C.P.B.) are gratefully acknowledged.

References

- [1] Racah, G., Lincei-Rend. Sc. fis. mat. e nat. **VIII**, 108-112 (1950) and Princeton lectures, CERN-61-8; Biedenharn, L. C., J. Math. Phys. **4**, 436-445 (1963); Klein, A., J. Math. Phys. **4**, 1283-1284 (1963); Gruber, B. and Raifeartaigh, O., J. Math. Phys. **5**, 1796-1804 (1964); Perelomov, A. M. and Popov, V. S., Math. USSR-Izvestija **2**, 1313-1335 (1968); Okubo, S., J. Math. Phys. **23**, 8-20 (1982)
- [2] For an outlook see R. Jackiw and B. Zumino in S. B. Treiman *et al.*, *Current algebra and anomalies*, World Sci. (1985). See also Kephart, T. W., Phys. Lett. **B151**, 267-270 (1985) and Okubo, S. and Patera, J., Phys. Rev. **D31**, 2669-2671 (1985). Further references may be found in the above articles and in de Azcárraga, J. A. and Izquierdo, J. M., *Lie groups, Lie algebras, cohomology and some applications in physics*, Camb. Univ. Press (1995)
- [3] de Azcárraga, J. A. and Pérez Bueno, J. C., *Higher-order simple Lie algebras* (hep-th/9605213), to appear in Commun. Math. Phys.
- [4] de Azcárraga, J. A.; Perelomov, A. M. and Pérez Bueno, J. C., J. Phys. **A29**, L151-L157 (1996); *The Schouten-Nijenhuis bracket, cohomology and generalized Poisson structures* (hep-th/9605067), J. Phys. **A**, in press, and these proceedings.
- [5] Nambu, Y., Phys. Rev. **D7**, 2405-2412 (1973); Takhtajan, L., Commun. Math. Phys. **160**, 295-315 (1994)
- [6] Lada, T. and Stasheff, J., Int. J. Theor. Phys. **32**, 1087-1103 (1993)